

A Novel Variant of N-R Method and its Convergence

Bhagwati Prasad and Ritu Sahni

Department of Mathematics
Jaypee Institute of Information technology,
A-10, Sector-62, Noida(UP) 201307 India

Copyright © 2014 Bhagwati Prasad and Ritu Sahni. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The intent of this paper is to propose a new iterative method from the well known Newton-Raphson (N-R) method for solving nonlinear equations. We also study the error estimate and the rate of convergence for the same.

Keywords: Newton-Raphson method, iterative methods, Lagrange multiplier method, order of convergence.

1. Introduction

The equation $f(x)=0$ can be rearranged in the form of a fixed point equation $g(x)=f(x)+x=x$. The solution(s) to such type of equation is(are) computed iteratively through some iterative procedure. The Newton-Raphson ((N-R) method is one of the most popularly used iterative methods for solving such nonlinear equations. If x_{n+1} represents the $(n+1)^{\text{th}}$ iterate of the N-R method, then we have

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (1)$$

The N-R method and other iterative methods are extensively studied by various authors and a number of variants of them appeared in the literature, see for instance, [1-3], [7-8], [10-15] and references therein. In this paper, we propose a new iterative algorithm from the N-R method and study the error estimate and rate of convergence of it.

2. New iterative Algorithm

We will use the general Lagrange multiplier method to re-derive the N-R method. If x_n is an approximate root of $f(x) = 0$, then $f(x_n) \neq 0$. We may write a correction equation in the following form

$$x_{n+1} = x_n + \lambda f(x_n), \quad (2)$$

where λ is a general Lagrange multiplier [6], which can be identified optimally by setting

$$\frac{dx_{n+1}}{dx_n} = 0 \quad (3)$$

We, therefore, can identify the multiplier as follows

$$\lambda = -\frac{1}{f'(x_n)} \quad (4)$$

This on substituting into (2) gives the well-known Newton-Raphson formulation. The above idea was first proposed by Inokuti, and was further developed to the well-known variational iteration method (see [4-5] and [9]). The correction equation (2) may be re-written in a more general manner as follows:

$$x_{n+1} = x_n + \lambda (1 + x_n)^\alpha f(x_n), \quad (5)$$

where α is a free parameter. Setting $\frac{dx_{n+1}}{dx_n} = 0$, we obtain

$$\lambda = -\frac{1}{\alpha(1 + x_n)^{\alpha-1} f(x_n) + (1 + x_n)^\alpha f'(x_n)}. \quad (6)$$

Hence the iteration formulation becomes

$$x_{n+1} = x_n - \frac{(1 + x_n)}{\alpha f(x_n) + (1 + x_n) f'(x_n)} f(x_n). \quad (7)$$

The free parameter α can be used as a control parameter to adjust convergence in every step if necessary. The optimal value for α can be determined from the relation $\frac{\partial x_{n+1}}{\partial \alpha} = 0$.

3. Numerical Examples

To illustrate the effectiveness of the iterative scheme (7), we consider the following examples:

Example 1. $x^{10} - 1 = 0$, $x_0 = 0.5$

Example 2. $e^{\sin x} - x = 0$, $x_0 = 0$

Example 3 (a). $x \sin x + \cos x = 0$, $x_0 = 1.0$ (b). $x \sin x + \cos x = 0$, $x_0 = 0$

Example 4. $x^3 - 3x + 1 = 0$, $x_0 = 1.0$

Example 5. $x - \cos(x) = 0$, $x_0 = 4.0$

Example 6. $\sin(x) = 0$, $x_0 = 1.6$.

The following table illustrates the nature of solutions of these examples at different number of iterations.

Example	New Iterative Algorithm			Newton- Raphson Algorithm	
	α	Number of iterations	x_n	Number of iterations	x_n
1	-2	6	1.0000	42	1.0000
2	-2	3	2.2191	20	2.2191
3(a)	-2	4	2.7984	6	56.531
3(b)	-2	5	2.7984	-----	Divergent
4	-2	4	1.5321	-----	Divergent
5	1.5	5	0.7391	28	0.7391
6	-2	4	3.1416	7	31.4159

Table 1. Comparison between New Iterative Algorithm and N-R method

It is clear from Table 1 that the proposed scheme converges faster than N-R method and thus it is more effective in many cases.

4. Rate of convergence

Definition 4.1. If a sequence $\{x_k\}$ converges to a root R then we say it converges

linearly to R if $\lim_{k \rightarrow \infty} \frac{|x_{k+1} - R|}{|x_k - R|} = \mu$, where $0 < \mu < 1$.

Definition 4.2. If a sequence converges quadratically to a root R then

$\lim_{k \rightarrow \infty} \frac{|x_{k+1} - R|}{|x_k - R|^2} = \mu$, where $0 < \mu < 1$.

Lemma 4.1. If a root x_0 has multiplicity $k > 1$ and the orbit of the new iterative scheme (7) converges, then the orbit converges linearly.

Theorem 4.1. If a root x_0 has multiplicity 1 and the orbit of the new iterative scheme (7) converges, then the orbit converges quadratically.

Proof. To prove that the scheme (7) converges quadratically for a root of multiplicity 1, we first express $\lim_{k \rightarrow \infty} \frac{|x_{k+1} - R|}{|x_k - R|^2}$ as $\lim_{k \rightarrow \infty} \frac{|E_{k+1}|}{|E_k|^2}$, where E represents the error term. Consider the Taylor polynomial of a function $f(x)$ whose roots we wish to compute around the point x_k . Assume that $f'(x_k) \neq 0$, then

$$f(x) = f(x_k) + f'(x_k)(x - x_k) + \frac{f''(x_k)(x - x_k)^2}{2} \quad (8)$$

If $f(x)$ has a root at $x = x_r$, then from above, we easily get

$$\frac{f'(x_k)}{f(x_k)} = \frac{-f''(x_k)(x_r - x_k)}{2} - \frac{1}{x_r - x_k}$$

Now the iteration formula (7) can be written as

$$\frac{f'(x_k)}{f(x_k)} = \frac{-1}{x_{k+1} - x_k} - \frac{\alpha}{1 + x_k} \quad (9)$$

and thus we get $\frac{E_{k+1}}{E_k^2 - E_k E_{k+1}} = \frac{1}{2} \frac{E_k f''(x_k)}{f(x_k)} - \frac{\alpha}{1 + x_k}$.

This on multiplying by $f(x_k)$ and simplifying gives

$$\frac{E_{k+1}}{E_k^2} = \frac{1}{2} f''(x_k)(x_{k+1} - x_k) - \frac{\alpha f(x_k)(x_{k+1} - x_k)}{(1 + x_k)(x_r - x_k)} \quad (10)$$

If, in fact, the orbit does converge to a root then $\lim_{k \rightarrow \infty} f'(x_k) = f'(x_r)$ and

$\lim_{k \rightarrow \infty} f''(x_k) = f''(x_r)$, so that $\lim_{k \rightarrow \infty} \frac{|E_{k+1}|}{|E_k|^2} = \frac{1}{2} |f''(x_k)(x_{k+1} - x_k)| > 0$.

Notice that $f'(x_r) \neq 0$, because x_r has multiplicity 1. If x_r has multiplicity $k > 1$ then, applying Lemma 4.1, $f(x)$ can be written as $f(x) = (x - x_r)^k G(x)$. Therefore, $f'(x) = k(x - x_r)^{k-1} G(x) + (x - x_r)^k G'(x) \Rightarrow f'(x_r) = 0$.

Acknowledgements. The authors would like to thank the referees.

References

1. C. Chun, Construction of Newton-like iteration methods for solving nonlinear equations, *Numer. Math.*, 104 (3) (2006), 297-315.
2. J. H. He, A modified Newton-Raphson method, *Commun. Numer. Methods Eng.* 20 (2004), 801-805.

3. J. H. He, Improvement of Newton iteration method, *Int. J. Nonlinear. Sci. Numer. Simulation*, 1(2) (2000), 239-240.
4. J. H. He, Some asymptotic methods for strongly nonlinear equations, *Int. J. Mod. Phys. B*, 20 (2006), 1141-1199.
5. J. H. He, Variational iteration method-a kind on non-linear analytical technique: some examples, *Int. J. Non-Linear Mech.*, 34(4) (1999), 699-708.
6. M. Inokuti, H. Sekine, and T. Mura, General use of the Lagrange multiplier in nonlinear mathematical physics, Variational method in the mechanics of solids. In: Nemat-Nasser S (ed.) (1978), 156-162.
7. A. B. Israel, Newton's method with modified functions, *Contemporary Mathematics* 204 (1997), 39-50.
8. V. Kanwar and Tomar S. K. Tomar, Modified families of Newton, Halley and Chebyshev methods, *Appl. Math. Comput.* 192(1) (2007), 20-26.
9. Z. M. Odibat and S. Momani, Application of variational iteration method to Nonlinear differential equations of fractional order *Int. J. Nonlinear Sci.* 7(2006) 27-34.
10. B. Prasad and R. Sahni, Stability of a general iterative algorithm, *Selected Topics in Applied Computer Science* (2010), 216-221.
11. B. Prasad and R. Sahni, A new method for solving nonlinear equations, *World Academy of Science, Engineering and Technology* 7(75) (2011), 599-604.
12. B. Prasad and R. Sahni, Convergence of some general iterative schemes, *Int. Journal of Math. Analysis* 5 (25) (2011), 1237 – 1242.
13. B. Prasad and R. Sahni, A convergence theorem for Jungck-Ishikawa iteration", *Recent Researches in Artificial Intelligence, Knowledge Engineering and Data Bases* (2011), 79-84.
14. W. X. Qian, Y. H. Ye, J. Chen, L. F. Mo, He's iteration formulation for solving nonlinear algebraic equations, *International symposium on nonlinear dynamics* (2007), 1-6.
15. J. R. Sharma, A composite third order Newton-Steffensen method for solving nonlinear equations, *Appl. Math. Comput.* 169(1) (2005), 242-246.

Received: January 11, 2014